

The Blazhko behaviour of RR Geminorum I

CCD photometric results in 2004

J. Jurcsik¹, Á. Sódor², M. Váradi^{1,3}, B. Szeidl¹, A. Washuettl⁴, M. Weber⁴, I. Dékány², Zs. Hurta²,
B. Lakatos², K. Posztobányi², A. Szing², and K. Vida²

¹ Konkoly Observatory of the Hungarian Academy of Sciences. P.O. Box 67, H-1525 Budapest, Hungary
e-mail: jurcsik@konkoly.hu

² Eötvös Loránd University, Department of Astronomy, P.O. Box 32, H-1518 Budapest, Hungary

³ Department of Experimental Physics and Astronomical Observatory, University of Szeged, H-6720 Szeged, Dóm tér 9, Hungary

⁴ Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

Received ; accepted

Abstract. Extended CCD monitoring of RR Gem revealed that it is a Blazhko type RRab star with the shortest Blazhko period (7^d.23) and smallest modulation amplitude ($\Delta M_{max} < 0.1$ mag) currently known. The short period of the modulation cycle enabled us to obtain complete phase coverage of the pulsation at each phase of the modulation. This is the first multicolour observation of a Blazhko star which is extended enough to define accurate mean magnitudes and colours of the variable at different Blazhko phases. Small, but real, changes in the intensity mean colours at different Blazhko phases have been detected. The Fourier analysis of the light curves shows that, in spite of the mmag and smaller order of the amplitudes, the triplet structure is noticeable up to about the 14th harmonic. The modulation is concentrated to a very narrow, 0.2 phase range of the pulsation, centred on the supposed onset of the H emission during rising light. These observational results raise further complications for theoretical explanation of the long known but poorly understood Blazhko phenomenon.

Key words. Stars: individual: RR Gem – Stars: variables: RR Lyr – Stars: oscillations – Stars: horizontal-branch – Techniques: photometric –

1. Introduction

Most RR Lyrae stars repeat their light curves with remarkable regularity. About 30% of the known galactic RR Lyrae stars, however, display cyclic modulation in the shape and amplitude of the light curve over tens to thousands of pulsational cycles (Smith 1995). The phenomenon is often called the Blazhko effect since Blazhko (1907) was the first to recognize that the light maximum of an RR Lyrae star (namely RW Dra) showed phase modulation on a long time-scale (around 42 days).

In several well-studied RR Lyrae stars, the amplitude and the period of the Blazhko effect has been observed to be variable on a time-scale of years (Szeidl 1988; Smith et al. 1999; LaCluyzé et al. 2004). In this respect RR Geminorum provides a very interesting case. Detre (1970) remarked that his unpublished photographic observations from the 1930s demonstrated the Blazhko effect in RR Gem, but (in his words) it ‘ceased to exist since about 1940.’ That the Blazhko effect might be temporary in nature may be crucial in understanding the physical

background of the phenomenon. Therefore we decided to begin an intensive photometric investigation of the star.

RR Gem ($\alpha = 07^h 21^m 33^s.53$, $\delta = +30^\circ 52' 59''.5$, J2000) was discovered by L. P. Ceraski (Ceraski 1903) on photographic plates taken by Blazhko. The first visual observations of the star was made by Graff (1905) and he was able to determine the pulsation period (0^d.397). Since that time a great number of visual observations have been published, but there are only a few published photoelectric observations available (e.g., Fitch et al. 1966; Liu & Janes 1989).

The absolute parameters of RR Gem were determined by Liu & Janes (1990) from a Baade-Wesselink analysis. It belongs to the metal rich ($[Fe/H] = -0.14$ dex as compiled in Jurcsik & Kovács (1996)) disk population group of RR Lyrae stars (Layden 1995).

Most of the papers dealing with Blazhko stars list the possible theoretical explanations of the phenomena and also the different observational arguments (see e.g. LaCluyzé et al. 2004; Alcock et al. 2003; Chadid et al. 2004). Neither of the two groups of possible models




Fig. 1. Folded V light curve of RR Gem. The scatter of the light curve is caused by Blazhko modulation and not by observational inaccuracies. The amplitude of the modulation is very small, it does not reach 0.1 mag even in maximum light, and the shape of the modulation points to a very regular behaviour. In this, and in the other figures, phase 0.5 is set to the middle of the ascending branch defined as the phase where the V flux is equal to its time averaged value.

a) the magnetic oblique rotator (Shibahashi 2000; Cousens 1983)

b) resonant coupling of nonradial mode(s) to the dominant radial mode (Nowakowski & Dziembowski 2001) can fully explain the large variety of observational results of Blazhko behaviour. It is important to note, however, that both of the models connect the Blazhko period to the rotation of the star.

In this first part of a series of two papers on RR Gem, we examine the photometric behaviour of the star exclusively using recent CCD observations. A reexamination of all the previous observations in order to reveal the long term behaviour of the modulation will be the subject of the second paper.

2. The data

The observations were obtained with the automated 60cm telescope of Konkoly Observatory, Svábhegy, Budapest equipped with a Wright 750x1100 CCD (parameters and calibration are given in Bakos (1999)) using $BV(RI)_c$ filters. The field of view was $17' \times 24'$. Data reduction was performed using standard IRAF¹ packages. More than 3000 frames were obtained in each passband on 56 nights between 14 January and 4 May in 2004 (JD 2453 019 – 2453 130). The data were transformed to the standard

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.




Fig. 2. Amplitude spectrum of the V light curve (top panel). Middle panel shows the residual spectrum after prewhitening with the pulsation frequency and its harmonics (kf_0 , $k = 1...15$). The location of the modulation frequency (f_m) and the symmetrically placed modulation frequency side lobes ($kf_0 \pm f_m$) are indicated by arrows. The spectral window is shown in the bottom panel.

system by using the U, B, V, R, I magnitudes given by Liu & Janes (1989) for SAO 060004 and SAO 060008. Relative aperture photometry to BD +31°1549 was calculated, and SAO 060004 was used as check star to control the stability of the results. The magnitude differences between our comparison and check stars remained constant with 0.0081, 0.0089, 0.0095, and 0.0093 mag *r.m.s.* scatter in the B, V, R, I colours, respectively.

The magnitudes and colours we derived for BD +31°1549 ($V = 10.277$ mag, $B - V = 0.366$ mag, $V - R = 0.233$ mag, and $V - I = 0.448$ mag) are in excellent agreement with the $V = 10.273$ mag and $B - V = 0.370$ mag values given by Sturch (1966). Photometric data are available electronically at the CDS. In the electronic tables (<http://cdsweb.u-strasbg.fr/cats/J.A+A.htx....>) Column 1 lists the HJD of the observation, and Column 2 gives the differential magnitude of RR Gem with respect to BD +31°1549 for the B, V, R, I colours.

3. Photometric results

The CCD observations revealed that, indeed, RR Gem shows Blazhko modulation, however, with very small, ~ 0.1 mag modulation amplitude. The modulation pattern is very regular, with a definite node (fixed point) on the ascending branch. Fig. 1 shows the V band observations superimposed on the mean light curve. The variation in the heights of maxima indicated a modulation period of about 7 days.

The Fourier spectra of the V data and the data prewhitened with the pulsation frequency and its harmonics are shown in Fig. 2. Data analysis was performed

Table 1. Amplitudes and phases of the pulsation and modulation frequency components of RR Gem.

		<i>B</i>		<i>V</i>		<i>R</i>		<i>I</i>	
A_0 / rms [mag]		10.822	0.013	11.418	0.011	11.152	0.013	10.906	0.012
f		A	ϕ	A	ϕ	A	ϕ	A	ϕ
[c/d]		[mag]	[deg]	[mag]	[deg]	[mag]	[deg]	[mag]	[deg]
f_0	2.517063	0.5527	80.0	0.4009	77.1	0.3080	72.9	0.2376	66.7
$2f_0$	5.034126	0.3111	296.1	0.2303	295.8	0.1799	294.7	0.1395	293.2
$3f_0$	7.551189	0.1751	169.7	0.1327	169.9	0.1055	169.6	0.0828	169.5
$4f_0$	10.068252	0.1184	46.2	0.0900	46.1	0.0707	45.6	0.0557	45.9
$5f_0$	12.585315	0.0663	278.0	0.0509	277.7	0.0409	276.4	0.0322	277.8
$6f_0$	15.102378	0.0494	142.9	0.0385	142.9	0.0304	143.1	0.0241	142.2
$7f_0$	17.619441	0.0341	18.3	0.0265	18.9	0.0212	18.7	0.0165	18.3
$8f_0$	20.136504	0.0217	239.5	0.0170	241.4	0.0133	241.5	0.0099	241.8
$9f_0$	22.653567	0.0174	105.1	0.0134	108.9	0.0109	109.5	0.0086	109.2
$10f_0$	25.170630	0.0125	333.2	0.0098	335.4	0.0071	338.7	0.0057	330.0
$11f_0$	27.687693	0.0091	196.2	0.0074	198.9	0.0054	195.5	0.0034	192.9
$12f_0$	30.204756	0.0072	57.2	0.0048	58.34	0.0044	58.6	0.0030	61.0
$13f_0$	32.721819	0.0055	282.0	0.0042	280.3	0.0036	294.2	0.0021	285.9
$14f_0$	35.238882	0.0044	145.8	0.0031	148.2	0.0026	141.3	0.0015	149.1
$15f_0$	37.755945	0.0039	3.9	0.0027	1.2	0.0024	2.7	0.0013	359.1
$f_0 - f_m$	2.378743	0.0093	185.6	0.0065	186.6	0.0059	196.2	0.0044	192.6
$f_0 + f_m$	2.655383	0.0092	349.2	0.0064	347.3	0.0058	350.9	0.0039	352.4
$2f_0 - f_m$	4.895806	0.0092	29.3	0.0073	38.8	0.0052	32.6	0.0044	42.7
$2f_0 + f_m$	5.172446	0.0086	175.3	0.0060	179.1	0.0043	181.2	0.0039	186.5
$3f_0 - f_m$	7.412869	0.0078	248.6	0.0061	248.4	0.0052	262.8	0.0038	271.1
$3f_0 + f_m$	7.689509	0.0071	43.9	0.0052	43.7	0.0046	45.8	0.0036	54.4
$4f_0 - f_m$	9.929932	0.0070	155.3	0.0054	156.3	0.0040	149.5	0.0032	161.5
$4f_0 + f_m$	10.206572	0.0067	306.8	0.0048	302.8	0.0043	304.1	0.0035	303.5
$5f_0 - f_m$	12.446995	0.0056	18.4	0.0043	14.8	0.0034	16.3	0.0028	16.4
$5f_0 + f_m$	12.723635	0.0054	190.0	0.0042	185.6	0.0033	188.6	0.0030	183.6
$6f_0 - f_m$	14.964058	0.0060	253.1	0.0043	250.4	0.0034	255.9	0.0027	243.3
$6f_0 + f_m$	15.240698	0.0035	41.3	0.0037	40.1	0.0024	38.1	0.0018	56.0
$7f_0 - f_m$	17.481121	0.0049	135.6	0.0036	136.1	0.0029	128.2	0.0027	126.9
$7f_0 + f_m$	17.757761	0.0048	280.5	0.0040	279.3	0.0037	282.5	0.0021	279.7
$8f_0 - f_m$	19.998184	0.0033	358.2	0.0026	1.2	0.0020	4.0	0.0026	356.5
$8f_0 + f_m$	20.274824	0.0033	169.6	0.0027	168.2	0.0018	172.1	0.0018	174.1
$9f_0 - f_m$	22.515247	0.0036	245.5	0.0025	240.8	0.0031	242.6	0.0017	246.8
$9f_0 + f_m$	22.791887	0.0030	30.6	0.0020	29.8	0.0017	21.4	0.0010	34.8
$10f_0 - f_m$	25.032310	0.0025	114.8	0.0020	100.3	0.0018	130.7	0.0014	113.8
$10f_0 + f_m$	25.308950	0.0024	269.4	0.0025	263.3	0.0012	254.8	0.0008	288.3
$11f_0 - f_m$	27.549373	0.0020	332.5	0.0017	339.3	0.0014	2.6	0.0014	325.4
$11f_0 + f_m$	27.826013	0.0018	138.7	0.0012	141.8	0.0012	152.6	0.0009	150.7
$12f_0 - f_m$	30.066436	0.0019	205.0	0.0009	205.5	0.0014	223.3	0.0011	206.3
$12f_0 + f_m$	30.343076	0.0016	356.0	0.0008	8.6	0.0007	343.0	0.0013	19.0
$13f_0 - f_m$	32.583499	0.0020	115.9	0.0015	111.0	0.0010	103.8	0.0003	147.5
$13f_0 + f_m$	32.860139	0.0019	252.5	0.0013	237.3	0.0006	223.0	0.0010	255.9
$14f_0 - f_m$	35.100562	0.0008	344.8	0.0008	330.2	0.0009	315.7	0.0006	329.6
$14f_0 + f_m$	35.377202	0.0011	155.2	0.0009	159.8	0.0013	122.4	0.0009	101.7
$15f_0 - f_m$	37.617625	0.0005	161.0	0.0007	219.6	0.0006	178.5	0.0006	140.6
$15f_0 + f_m$	37.894265	0.0008	11.8	0.0004	354.5	0.0005	30.4	0.0002	155.3
f_m	0.138320	0.0058	345.9	0.0045	355.7	0.0040	345.9	0.0042	354.1

using the different applications of the MUFAN package (Kolláth 1990). The Fourier spectrum of the residual shows the triplet structure typical of amplitude modulation up to about the 14th harmonic component. The side lobe frequencies have similar amplitudes and are symmetrically placed around the radial mode frequency and

its harmonics. The amplitudes of the side lobes decrease much less steeply than the amplitudes of the harmonics of the main pulsation frequency (see Fig. 3). This fact may imply a different explanation of these frequency components than that of the radial mode where nonlinear effects invoke harmonic components with exponentially decreas-

Table 2. Side lobe amplitude ratios, phase differences, and their errors. $R_k = A_{kf_0+f_m}/A_{kf_0-f_m}$; $\Delta\phi_k = \phi_{kf_0+f_m} - \phi_{kf_0-f_m}$.

	<i>B</i>				<i>V</i>				<i>R</i>				<i>I</i>			
<i>k</i>	R_k	σ_{R_k}	$\Delta\phi_k$	$\sigma_{\Delta\phi_k}$	R_k	σ_{R_k}	$\Delta\phi_k$	$\sigma_{\Delta\phi_k}$	R_k	σ_{R_k}	$\Delta\phi_k$	$\sigma_{\Delta\phi_k}$	R_k	σ_{R_k}	$\Delta\phi_k$	$\sigma_{\Delta\phi_k}$
1	0.989	0.045	163.6	3.0	0.970	0.063	160.7	3.7	0.983	0.071	154.7	4.9	0.867	0.088	159.8	6.2
2	0.934	0.053	146.0	3.3	0.822	0.053	140.3	3.8	0.827	0.100	148.6	6.3	0.886	0.091	143.8	6.5
3	0.910	0.060	155.3	3.7	0.852	0.065	155.3	4.1	0.885	0.103	143.0	5.7	0.947	0.109	143.3	6.8
4	0.957	0.070	151.5	3.9	0.887	0.076	146.5	4.6	1.075	0.128	154.6	6.6	1.063	0.137	142.0	7.4
5	0.982	0.088	171.6	5.2	0.977	0.098	170.8	5.7	0.971	0.144	172.3	8.8	1.034	0.176	167.2	9.0
6	0.583	0.077	148.2	6.5	0.860	0.092	149.7	5.9	0.706	0.144	142.2	10.3	0.667	0.134	172.7	11.8
7	0.980	0.114	144.9	6.0	1.111	0.125	143.2	6.5	1.276	0.224	154.3	9.1	0.778	0.141	152.8	11.5
8	1.000	0.129	171.4	8.6	1.038	0.166	167.0	9.2	0.900	0.269	168.1	15.5	0.692	0.140	177.6	12.7
9	0.833	0.108	145.1	8.9	0.800	0.154	149.0	11.2	0.548	0.110	138.8	13.8	0.588	0.205	148.0	22.5
10	1.000	0.170	154.6	11.4	1.250	0.240	163.0	10.7	0.632	0.187	124.1	20.6	0.714	0.263	174.5	24.0




Fig. 3. Amplitude ratios of the harmonic components of the dominant radial pulsation mode A_{kf_0}/A_{f_0} (solid) compared to the amplitude ratio behaviour of the modulation components $A_{kf_0-f_m}/A_{f_0-f_m}$, $A_{kf_0+f_m}/A_{f_0+f_m}$ (open symbols). While the amplitudes of the pulsation decrease exponentially, the amplitude decrease of the modulation is linear. Error bars for the modulation components' amplitude ratios are indicated, the errors of the amplitude ratios of the dominant pulsation mode are smaller than the symbols used.

ing amplitudes. The modulation components' amplitudes of AR Her (a Blazhko variable with large amplitude and phase modulation and a 32 d modulation period) show similar behaviour; from the fourth harmonic component its modulation amplitudes dominate over the pulsation ones (Smith et al. 1999).

Table 1 summarizes the amplitudes and phases of the pulsation and modulation frequencies in the *B*, *V*, *R*, *I* passbands calculated from the least squares solution using sine terms and HJD $T_0 = 2453019.0$ initial epoch value according to the formula given in Kovács (1995, Eq. 2). In the top of Table 1 the mean magnitudes (A_0) and the standard deviations of the fits are also given.

This solution has been calculated by using locked values of both the pulsation and modulation frequency components. The pulsation frequency adopted is the mean value obtained for the *B*, *V*, *R*, *I* data. The modulation frequency has been determined as the frequency which results the smallest *r.m.s.* scatter of the residuals of the least squares solutions assuming symmetrical displacement of the side lobes up to the 9th harmonic component of the radial mode. The period of the modulation obtained this way is 7^d23. In Fig. 4 the *V* light curve is folded according to this modulation period; sinusoidal variation in the height of maximum light is evident. The ephemerides of maximum pulsation light and maximum pulsation amplitude determined from the measurements of the 2004 observing season are the followings:

$$\text{HJD } T_{max} = 2453019.1590 + 0.3972884 \cdot E_{\text{pulsation}}$$

$$\text{HJD } T_{Blmax} = 2453021.30 + 7.23 \cdot E_{\text{Blazhko}}.$$

The error weighted mean value of the asymmetry parameter ($Q = \frac{A_+ - A_-}{A_+ + A_-}$) averaged over the first 10 modulation component pairs in the four colours is -0.035 ± 0.057 , indicating very small asymmetry with slightly larger amplitudes at the shorter frequency sides. For comparison the distribution of this parameter of the MACHO data peaks at +0.3 (Alcock et al. 2003).




Fig. 4. The *V* light curve of RR Gem folded with the 7^d23 Blazhko period. The heights of maximum light clearly show a 0.1 mag amplitude modulation.




Fig. 5. *V* data folded with the Blazhko period (left panels) and their residual spectra after prewhitening with the pulsation period and its harmonics (right panels) for two equal length parts of the dataset. In the first half of the observations (between JD 2453019 and 2453174) the data coverage is nearly the same as for the whole data set and the residual spectrum is very similar to that shown in Fig. 2 for the complete data set (top panels). In the second part of the observations (between JD 2453077 and 2453130) the data are more sparse, and the residual spectrum has changed significantly. The near equality of the sidelobe amplitudes has disappeared, instead, highly asymmetrical structure emerge, with very small if any power at the higher frequency sides (bottom panels).

The currently viable theoretical explanations of the Blazhko phenomenon predict approximate equality of the amplitude ratios, and phase differences between the side lobe frequencies of the different order components in each colour (Kovács 1995; Shibahashi 2000; Smith et al. 1999; Nowakowski & Dziembowski 2001). Although these predictions are based on simplified models it is worth checking the stability of these components in our data. Table 2 lists the amplitude ratios and phase differences and their errors of the side lobe frequency pairs in the *B*, *V*, *R*, *I* bandpasses for the first 10 modulation component pairs. Differences larger than 2σ can be found in the amplitude ratios while the phase differences of the first 5 modulation component pairs in the different colours do not agree within the 3σ limit. The mean values and the standard deviations of the phase differences in the *B*, *V*, *R*, *I* colours of the first five modulation component pairs are 159.7 ± 3.7 , 144.7 ± 3.5 , 149.2 ± 7.0 , 148.7 ± 5.5 , and 170.5 ± 2.3 . The phase differences of the fifth components are statistically significantly larger in each colour than the phase differences of the second pairs. The large scatter of the third and fourth components indicates colour to colour changes of the phase differences, too. Differences between these parameters (R_k , $\Delta\phi_k$) have also been detected in AR Her (Smith et al. 1999).

3.1. Stability of the results

One of the theoretically unexplained characteristics of Blazhko stars is the high asymmetry of the amplitudes of the $\pm$ modulation frequency components. In contrast, only very small if any asymmetry can be seen in Fig. 2. We have checked what happens if data subsets with less complete phase coverage are used. The results are shown in Fig. 5. The 110 day dataset has been divided into two intervals of equal length but different number of observations (~ 2000 and ~ 1000). In the first part of the observations both the clear sky statistics were better, and the observational runs were longer. The residual spectrum of the first dataset is very similar to that of the complete dataset shown in the middle panel of Fig. 2. However, the residual spectrum of the second dataset is significantly different, with enhanced power at the lower frequency side of the modulation, and with diminishing amplitudes at the higher frequency side. The data distribution of this second dataset also contains observations from each 0.1 phase intervals of the Blazhko period, but for some phase intervals no complete light curve can be constructed. This is the typical data distribution of Blazhko star photometry, thus our result warns that classification schemes of the modulation based on the asymmetry of the side lobe amplitudes might be seriously biased by the phase coverage of the data and may lead to an over-interpretation of

the results. As the Blazhko period of RR Gem is about an order of magnitude shorter than the typical period of the modulation, it should be expected that for stars with longer modulation periods even more extended observations than we have on RR Gem are needed to correctly describe the properties of the modulation. For comparison, in Alcock et al. (2003), 700 – 1000 measurements of the stars were analysed.

The effect of data distribution biases on the side lobe amplitudes, however, has to be carefully treated. The fact that the different observations of RR Lyrae (Szeidl & Kolláth 2000; Smith et al. 2003) equally show the longer frequency modulation components to have higher amplitudes points to real differences in the amplitudes of the modulation components.

4. Variations during the Blazhko cycle

The full coverage of the pulsation light curve at different phases of the Blazhko cycle makes it possible to study the changes of the light curves' shapes and Fourier parameters in detail, and to detect any possible variation in the mean magnitudes and colours.

4.1. Light curve changes

We have divided the data into 10 subsets containing observations from different phases of the Blazhko cycle. The minimum and maximum number of data in these subsets are 165 and 479, respectively. The behaviour of the Fourier parameters of the light curves in the different Blazhko phases is shown in Fig. 6. The total range of the variations in the A_1 , A_{10} amplitudes are 0.026, (between 0.390 and 0.416 mag) and 0.009 mag (between 0.005 and 0.014 mag), respectively. The amplitude ratios ($R_{k1} = A_k/A_1$) follow the changes of the amplitudes but there is no significant change in the epoch-independent phase differences ($\phi_{k1} = \phi_k - k\phi_1$) in accordance with the finding that no phase modulation in the light curve changes has been detected.

We have checked how the light curves at different Blazhko phases deviate from the mean light curve. Fig. 7 and 8 show the pulsation light curves at each one-tenth phase of the Blazhko cycle, and the residual curves after subtracting the mean light curve from the data subsets.

Due to the good coverage of the data, a least squares fit with the pulsation period and its harmonics to the entire dataset (as shown in Fig. 1) is a good representation of the mean light curve of the variable. The Fourier parameters of this 'average' solution agree within the limits of the uncertainties with the corresponding amplitudes and phases of the complete solution listed in Table 1.

The most striking property of the residual curves is that they show intense variations only in a narrow, ~ 0.20 phase interval of the pulsation. The residual curve of all the data (Fig. 9) shows that the distortion is symmetrical to phase 0.5 which is set to the phase of the middle of the rising branch as described in Sect. 3. A similar

residual plot has been shown by Lee & Schmidt (2001) for DR And with the conclusion that 'large residuals are obtained from data points on the ascending branch'. To analyse such a behaviour with the Fourier decomposition method might not be the most appropriate tool, as for a reliable model very high harmonic modulation frequencies are needed. This is reflected in the significantly different amplitude decrease behaviour of the modulation frequencies from the amplitude decrease of the harmonic components of the main pulsation (see Fig. 3). Although no complete fully nonlinear model calculation exists, Nowakowski & Dziembowski (2003) showed that in evolved stars the nonradial modes are highly nonlinear. However, even in the case of high nonlinearity it is hard to explain why the high (5 – 10) order nonlinear coupling terms have amplitudes commensurable with the first order ones.

It is also important to think about why the modulation is concentrated in the minimum to maximum phase interval of the pulsation, where nonlinear effects are most important. In the pioneering spectroscopic study of




Fig. 6. Fourier parameters of the V light curves in the different phases of the Blazhko cycle. The amplitude ratios vary in phase with the A_1 amplitude, but no changes in the epoch independent phase differences can be detected.




Fig. 7. Pulsation light curve constructed from data falling into the 0.1 phase intervals of the Blazhko cycle. In each plot the mean light curve is also shown. Slight, but systematic differences of the observations in most of the plots are evident.

RR Lyrae, Preston et al. (1965) concluded that a displacement of the shock-forming layer in the atmosphere takes place during the Blazhko cycle. In Preston & Paczyński (1964) it was also shown that: ‘the phases at which H emission occurs are closely related to phases of photometric parameters’. Thus the phase of the $U-B$ excess and the phase on the ascending branch when the visual luminosity of the star equals its time average value (at phase 0.5 in the figures) are closely related, with only some minute differences, to the onset of the H emission. The $U-B$ observations of Liu & Janes (1989) indicate strong excess around the phase of the mid ascending branch in RR Gem. In Figs. 8 and 9 the symmetrical modulation is centred exactly on this phase of the pulsation, indicating a connection between the origin of the modulation and that of the H emission.

As the standard deviations of the data from the fits using the 46 frequencies listed in Table 1 are larger in each colour than expected from the observations of the comparison and check stars, some further variations of the residual light curves are also suspected. The Fourier spectra of the residuals after prewhitening with both the pulsation and modulation frequencies listed in Table 1 for







Fig. 8. The same data as shown in Fig. 7 after subtracting the mean light curve. Intense variations are concentrated in a 0.20 phase interval centred on phase 0.5 which corresponds in our representation to the middle of the ascending branch, when H emission is supposed to occur.

the B, V, R, I data are shown in Fig. 10. Another modulation frequency component is present in each colour; the arrows indicate the positions of $f = kf_0 + f'_m$ ($k = 1..6$). This tertiary periodicity corresponds to modulation with a period of 6^d.69, which is very close to the dominant modulation period. A tertiary period shorter than the dominant modulation period has been also found by LaCluyzé et al. (2004) for XZ Cygni. A quintuplet structure with equidistant frequency spacing would predict secondary modulation with a much shorter, 3^d.61 periodicity.

Fig. 11 shows the residual V light curve after prewhitening with the frequencies of Table 1 folded with the pulsation period. Residuals are still concentrated in the same phase of the pulsation as in Fig. 9. This behaviour of the residuals indicates that, perhaps, some irregular behaviour of the Blazhko modulation mimics the tertiary periodicity. This idea seems to be supported by separate analyses of the two halves of the data, as neither parts of the data show the same residual structure shown in Fig. 11.

Table 3. Magnitude and intensity mean brightnesses and colours in the different Blazhko phases.

Bl phase	magnitude averages								intensity averages						
	$A_1(V)$	$\overline{B}$	$\overline{V}$	$\overline{R}$	$\overline{I}$	$\overline{B} - \overline{V}$	$\overline{V} - \overline{R}$	$\overline{V} - \overline{I}$	$\overline{B}$	$\overline{V}$	$\overline{R}$	$\overline{I}$	$\overline{B} - \overline{V}$	$\overline{V} - \overline{R}$	$\overline{V} - \overline{I}$
0.0 – 0.1	0.3898	11.819	11.414	11.150	10.903	0.405	0.264	0.511	11.713	11.359	11.118	10.885	0.354	0.241	0.474
0.1 – 0.2	0.3981	11.818	11.416	11.150	10.905	0.402	0.266	0.511	11.707	11.358	11.116	10.885	0.349	0.242	0.473
0.2 – 0.3	0.4036	11.824	11.419	11.153	10.908	0.405	0.266	0.511	11.706	11.358	11.117	10.888	0.348	0.241	0.470
0.3 – 0.4	0.4104	11.827	11.421	11.156	10.910	0.406	0.265	0.511	11.702	11.356	11.117	10.888	0.346	0.239	0.468
0.4 – 0.5	0.4159	11.829	11.422	11.154	10.911	0.407	0.268	0.511	11.701	11.355	11.115	10.888	0.346	0.240	0.467
0.5 – 0.6	0.4088	11.824	11.421	11.155	10.908	0.403	0.266	0.513	11.701	11.356	11.117	10.885	0.345	0.239	0.471
0.6 – 0.7	0.4081	11.824	11.420	11.154	10.908	0.404	0.266	0.512	11.704	11.357	11.117	10.886	0.347	0.240	0.471
0.7 – 0.8	0.4953	11.823	11.416	11.152	10.905	0.407	0.264	0.511	11.710	11.358	11.118	10.886	0.352	0.240	0.472
0.8 – 0.9	0.3947	11.820	11.414	11.150	10.903	0.406	0.264	0.511	11.712	11.357	11.116	10.883	0.355	0.241	0.474
0.9 – 1.0	0.3900	11.816	11.413	11.146	10.901	0.403	0.267	0.512	11.710	11.357	11.114	10.882	0.353	0.243	0.475

**Fig. 9.** The V residual light curve of RR Gem folded with the $0^d3972884$ pulsation period after subtracting the mean pulsation light curve.**Fig. 11.** The V residual light curve of RR Gem folded with the $0^d3972884$ pulsation period after prewhitening with the pulsation and modulation frequencies listed in Table 1. Residual light variation is still concentrated in the same $(0.4 - 0.6)$ phase interval of the pulsation as in Fig. 9.**Fig. 10.** Amplitude spectra of the residual B, V, R, I light curves after prewhitening with the pulsation and modulation frequencies listed in Table 1. Arrows indicate residual frequencies at $kf_0 + f'_m$ ($k = 1..6$).

4.2. Magnitude and colour changes

Multicolour photometry of Blazhko stars which were extended enough to determine whether there is any change in the mean colour of the variables during the Blazhko period has never been obtained.

According to the mean characteristics of the Blazhko variables in the MACHO database, the modulation frequency occurs in the spectra with an overall amplitude of 0.006 mag (Alcock et al. 2003), i.e., this is the typical amplitude of the modulation in the magnitude average V light level of Blazhko variables.

In Table 3 the magnitude and intensity mean B, V, R, I magnitudes of RR Gem in the different Blazhko phases are listed. Mean colours are calculated as the arithmetic differences between the mean magnitudes. Fig. 12 shows the mean $\overline{V}$ magnitude and $\overline{B} - \overline{V}$ and $\overline{V} - \overline{I}$ colour variations during the Blazhko cycle calculated as magnitude and in-

tensity averages. Although the magnitude mean brightness level of the star seems to vary with a 0.005 mag semi-amplitude (in accordance with the MACHO results), the intensity mean brightness shows much smaller if any variation. Concerning the colours, the situation is the opposite; the colours calculated from the intensity mean magnitudes definitely show correlated changes with the Blazhko phase, namely, the star is slightly cooler in its small amplitude phase than during the highest amplitude period.

Whether intensity or magnitude mean colors better correspond to the static values of pulsating stars is still controversial (Bono et al. 1995). As the luminosities of the stars have real physical meaning, we think that the intensity means reflect the real parameters of the star more correctly. According to the intensity mean values the mean brightness level of RR Gem remains the same during the Blazhko cycle, but a slight colour (temperature) variation takes place in the sense that the star is $\sim 30 - 40\text{K}$ cooler when the amplitude of the pulsation is the smallest.

5. Discussion

The extended CCD observations of RR Gem exposed some previously unknown or unexamined properties of the Blazhko phenomenon. The new results concerning the properties of the modulation of RR Gem are summarized in the next items.




Fig. 12. Intensity and magnitude mean $\bar{V}$ magnitude and $\bar{B} - \bar{V}$ and $\bar{V} - \bar{I}$ colours of RR Gem in the different Blazhko phases. The intensity mean colours indicate colour variations during the Blazhko cycle, namely the star is slightly cooler in the smaller amplitude phase than during the highest amplitude period.

- First detection of a very low amplitude Blazhko modulation with symmetrically placed triplets in the Fourier spectrum.

- The modulation is concentrated in a 0.2 phase interval of the pulsation, centred on the mid of the ascending branch.
- In contrast to the exponential decrease in the amplitudes of the harmonic components of the radial pulsation, the decrease in the amplitudes of modulating components is close to linear.
- In contrast to theoretical expectations the amplitude ratios and phase differences of the $\pm$ modulation components are not the same for the different order modulation component pairs and in the different colours.
- According to the intensity mean colours and magnitudes in the different phases of the Blazhko cycle, there is very small if any mean V brightness variation along the Blazhko period, while the mean colours indicate small but definite changes in the sense that the star is 30-40 K cooler in its small amplitude phase.
- The light curve cannot be completely fitted with symmetrically placed Fourier frequency components alone. Residual scatter of the light curve is still concentrated in the ascending branch, indicating some irregular behaviour of the modulation.

Any model which relates the Blazhko periodicity to the rotation of the star predicts a similar percentage of small amplitude modulation as large amplitude ones. Because of the incidental angle of view of the rotational axes, the observed amplitudes of the modulation has to vary uniformly from zero up to some tenths of a magnitude. On the contrary, no small amplitude Blazhko modulation has been known previously. This fact has to be carefully examined on the grounds of observational biases. The most extended observations of field RR Lyrae stars were the photographic observations from the middle of the 20th century. These observations were not, however, accurate enough to detect modulation amplitudes smaller than ~ 0.2 mag. Photoelectric observations of field RR Lyrae stars are very sparse and usually time limited, so again, they were not suitable to detect small amplitude, long period modulations. The cluster and extragalactic variable surveys are usually seriously affected by crowding problems and are also often very time limited, thus small amplitude modulations might be missed or attributed to observational inaccuracy.

The detection of hundredths of magnitude amplitude modulation of RR Gem indicates that there might be a serious selection effect against small amplitude Blazhko stars. A systematic search using today's accurate photometric detectors is needed to reveal the true unbiased statistics of the modulation amplitudes.

Among the 731 fundamental mode Blazhko variables of the MACHO database there was only one star with a modulation period shorter than 10 days and only 5 with $P_{Bl} < 15$ days (Alcock et al. 2003). In contrast, RR Gem is already the second very short modulation period variable in the much smaller sample of well observed galactic field Blazhko variables (less than 50 stars). The other short period Blazhko star (AH Cam) was studied in de-

tail by Smith et al. (1994). Both AH Cam and RR Gem are relatively metal rich variables with pulsation periods shorter than 0^d.4. These are their common features, but the modulation pattern of the two stars are very different. In AH Cam the amplitude of the modulation is 0.3 – 0.4 mag, and phase modulation with about 0.1 pulsation phases occurs as well. A simplified explanation of the larger percentage of short modulation period Blazhko stars in our Galaxy than in the LMC would relate the short modulation periods to the larger metal content of these stars. However, the 530 day modulation period of RS Boo (Kanyó (1986); Nagy (1998)) with $P_{puls} = 0.377$ d and $[Fe/H] = -0.35$ dex (Layden 1995) indicates that a more complex description is needed.

Though Chadid et al. (2004), favouring the nonradial mode explanation of the Blazhko phenomenon, argue against the presence of a strong magnetic field in RR Lyrae, our detailed analysis of RR Gem revealed a challenge to these models as well. The two most important results which would be hard to explain by nonradial modes are the followings.

a) The linear decrease of the amplitudes of the modulation components compared to the exponential decrease in the amplitudes of the radial mode (Fig. 3) indicates significant differences between these frequency components. As nonlinearity is proportional to the amplitude, it is expected that if the harmonic terms of the radial mode invoked by nonlinear coupling decrease exponentially, then the nonlinear coupling terms of the much smaller amplitude nonradial modes have to decrease even more steeply. The linear decrease of the amplitudes of the very small amplitude modulation components indicates a different origin of its higher order components than nonlinear coupling. This anomalous behavior of the amplitudes of the modulation frequencies arises from the high concentration of the modulation in only 0.2 phase intervals of the pulsation.

b) The modulation is centred on the phase when H emission is assumed to occur. Such a connection between a temporal physical phenomenon in the atmosphere and the occurrence of the modulation suggests that during the Blazhko period real changes in the atmosphere are present. This is also supported by the detected colour variations during the Blazhko cycle.

Our detailed photometric observations of RR Gem have given new insight into the Blazhko phenomenon. However, it is an open question whether the Blazhko properties of RR Gem are unique or give general information about the modulation as well. To find the answer, further similar studies of other Blazhko variables are planned.

Acknowledgements. This research has made use of the SIMBAD database, operated at CDS Strasbourg, France. The financial support of OTKA grants T-043504 and T-046207 is acknowledged.

References

- Alcock, C., Alves, D. R., Becker, A. et al. 2003, *ApJ*, 598, 597
- Bakos, G. Á. 1999, *Occ. Techn. Notes Konkoly Obs.*, No. 11, <http://www.konkoly.hu/mitteilungen/mitteilungen.html>
- Blazhko, S. 1907, *Astron. Nachr.*, 173, 325
- Bono, G., Caputo, F., & Stellingwerf R. F. 1995, *ApJS*, 99, 263
- Ceraski, W. 1903, *Astron. Nachr.*, 161, 363
- Chadid, M., Wade, G. A., Shorlin, S. L. S., & Landstreet, J. D. 2004, *A&A*, 413, 1087
- Cousens, A. 1983, *MNRAS*, 203, 1171
- Detre, L. 1970, *Transactions of the IAU XIV A. Reports on Astronomy*, 259
- Fitch, W. S., Wiśniewski, W. Z., & Johnson, H. L. 1966, *Comm. Lun. and Planet. Lab.*, Vol 5, Part 2, No 71
- Graff, K. 1905, *Mitteilungen d. Hamburger Sternwarte*, Bd.3, Nr.8
- Jurcsik, J., & Kovács, G. 1996, *A&A*, 312, 111
- Kanyó, S. 1986, *Mitt. der Sternw. der Ungarischen Akad. der Wiss.*, Nr. 87
- Kolláth, Z. 1990, *Occ. Techn. Notes Konkoly Obs.*, No. 1, <http://www.konkoly.hu/staff/kollath/mufran.html>
- Kovács, G. 1995, *A&A*, 295, 693
- LaCluyzé, A., Smith, H. A., Gill, E.-M., et al. 2004, *AJ*, 127, 1653
- Layden, A. C. 1995, *AJ*, 110, 2288
- Lee, K. M., & Schmidt, E. G. 2001, *PASP*, 113, 1140
- Liu, T., & Janes, K. A. 1989, *ApJS*, 69, 593
- Liu, T., & Janes, K. A. 1990, *ApJ*, 354, 273
- Nagy, A. 1998, *A&A*, 339, 440
- Nowakowski, R. M., & Dziembowski, W. A. 2001, *Acta Astron.*, 51, 5
- Nowakowski, R. M., & Dziembowski, W. A. 2003, *Ap&SS*, 284, 273
- Preston, G. W., & Paczyński, B. 1964, *ApJ*, 140, 181
- Preston, G. M., Smak, J., & Paczyński, B. 1965, *ApJS*, 12, 99
- Shibahashi, H. 2000, in *ASP Conf. Ser. 203, The Impact of Large-scale Surveys on Pulsating Star Research*, eds., L. Szabados and D.W. Kurtz, p. 299
- Smith, H. A., Matthews, J. M., Lee, K. M. et al. 1994, *AJ*, 107, 679
- Smith, H.A. 1995, *RR Lyrae Stars* (Cambridge University Press)
- Smith, H. A., Barnett, M., Silbermann, N. A., & Gay, P. 1999, *AJ*, 118, 572
- Smith, H. A., Church, J. A., Fournier, J. et al. 2003, *PASP*, 115, 43
- Sturch, C. 1966, *ApJ*, 143, 774
- Szeidl, B. 1988, in *Multimode Stellar Pulsation*, eds., G. Kovács, L. Szabados and B. Szeidl, (Kultúra, Budapest), p. 45
- Szeidl, B., & Kolláth, Z. 2000, *ASP Conf. Ser. Vol.203., RR Lyrae: Analysis of 100 years of Observations*, eds.: Szabados, L., Kurtz, D. W., p. 281